




Integrated Pest Management (IPM) Sub-Group 2020 Report

Coordinator: Anne Fisher, UK, USA SC
Liaison: Colin Fisher, UK, USA


CORESTA Congress Online, October 2020





Last Integrated Pest Management (IPM) SG Meeting

- ❖ **AP Meeting in Victoria Falls: October 2019**
 - **10 registered**
 - **24 attended**





Value of Previous IPM Work

❖ IPM is not new – INTEGRATED management system

➤ Zimbabwe, TRB handbook 1950's recommended

- Rotation for nematode control
- Hygiene for TMV control
- Avoiding over-fertilization for bacterial foliar disease control

➤ US grower guides 1940's recommended

- Rotation & hygiene for black shank control
- Hygiene for TMV control

❖ Some new IPM strategies

➤ Mostly built on well-established principles



How This Work Helps the Scientific Community

❖ Lower CPA residues – BIG issue for tobacco industry

- CPAs may be replaced or partly replaced by other strategies
- Lower levels applied
 - Scouting
 - Proper application
 - Less disease



❖ Lower diseases/pest populations – easier control, less CPAs

- Rotations, good hygiene etc.
 - Prevent or slow build-up of diseases / pests



❖ Sustainable production – soils, flora/fauna, disease/pest levels


- Will we be growing tobacco 20 years from now?

❖ 80 chapters over 5 groups

- diseases
- nematodes
- insects
- weeds
- IPM strategies

❖ Each with a group leader

- organizes group
- collects chapters
- arranges reviews



Same approach for 3 groups

Diseases



fungal
bacterial
viral
seedling
post-harv

Nematodes



Insects



- **Groups divided into sections**
- **Chapter for each disease or pest**

Weeds group

Field Weeds



Parasitic Weeds



Different approach

- Principles of weed control
- Specific weed problems

IPM Strategies

Biological Control



Rotation



Correct CPA Usage



➤ Sections deal with general IPM principles

Alternate Hosts

Many solanaceous weeds are hosts of this pathogen ([Ch. 61](#)). Examples are Apple of Peru (*Nicandra physaloides*) and Jimson weed / stinkblaar (*Datura stramonium*), shown in Fig.15.6. Such weeds should be removed from the proximity of the fields and especially seedbeds / greenhouses. This is particularly important in areas which do not have killing winter frosts, where weeds overwinter.

D.1. Field Weeds

61. Weeds as Alternate Hosts to Other Pests

Andy Bailey, University of Kentucky, USA

General

Weeds can act as a major host site for other tobacco pests such as diseases, nematodes, and insects. Many weeds that commonly occur around tobacco fields can harbor other pests and result in increased infection on tobacco crops. Generally, weed species that have the closest botanical relationship to tobacco, such as solanaceous weed species, are most likely to harbor pests that can infest tobacco. However, many plant species with little botanical relationship to tobacco can also serve as hosts.

- ❖ **Collect outstanding chapters**
 - **Some not done, some in progress**
 - Some new chapters received
 - New authors & leaders
- ❖ **Complete outstanding reviews, editing**
 - **Currently in progress**
 - **2 chapters ready for website review**
- ❖ **Document posted incomplete**
 - **Task Force → Sub-Group**
 - Add completed chapters
 - Update existing chapters



CORESTA
 International Centre for Scientific Research Relative to Tobacco
 Centre de Coopération pour les Recherches Scientifiques Relatives au Tabac

IPM Taskforce

Completed Documents

As each completed chapter has been reviewed and edited, it will be placed on this page for general review by the taskforce. Members will be notified by email when a document has been posted online. Please peruse any documents which are of interest to you, and let us know if you have any comments.

We would encourage those of you with experience of any particular topic to comment and contribute, especially if you know of any IPM strategy which has been omitted. We are particularly interested in a global perspective and welcome suggestions of different strategies from our diverse membership.

Each document will be posted online for one month. The post date and expiration date included in the table below will also be in the notification email. Please send comments to the author using the [Comments link](#) in the table; these comments will be automatically copied to the section leader (see [Taskforce Overview](#) page for assistance) and the [taskforce coordinator](#).

10/21/15: There are currently no chapters for website review, but several chapters will be posted shortly.

Document Name	Post Date	Expires	Author	Reviewer 1	Reviewer 2	*Comments: please contact

*Note: email comments to the author will automatically be carbon copied to the [IPM Taskforce coordinator](#) and to the relevant section leader.

[Return to Top](#) [Taskforce Overview](#) [Working Documents](#) [IPM Images](#) [Completed Documents](#)
[IPM Presentations](#) [Final Report](#) [Useful Links](#) [Contacts](#)
 Last Updated: October 22, 2015

[Return to Home Page](#)

❖ Our many members

- Photo contributors
- Reviewers
- Especially authors

❖ CORESTA



A.2. Bacterial Diseases

16. Wildfire, angular leaf spot *Pseudomonas syringae* pv. *tabaci* (tox+), (to-)
Anne Jack, University of Kentucky, USA

General

These diseases can affect tobacco in both the seedbeds / float trays and the field, although wildfire (tox+) tends to be more of a problem in the seedbed and angular leaf spot (tox-) in the field. Wildfire and angular leaf spot are not major problems in many tobacco producing areas, such as the USA, Brazil and Europe. In Africa, they are diseases of major importance which can cause devastating losses, especially in wet seasons. All control measures discussed refer only to areas where they are diseases of economic importance, and are not usually necessary in areas such as the USA.

Symptoms

The symptoms of the tox+ (toxin producing) and tox- forms of this disease are quite different. Wildfire (tox+) is characterized by a small brown or black water-soaked lesion, surrounded by a chlorotic halo (Figure 16.1). Wildfire can be systemic in seedlings, causing distortion (Figure 16.4). The angular (tox-) lesion is brown, dark brown or black, much larger than the wildfire lesion, has little or no chlorotic halo, and has angular margins because the lesion is confined by the lateral veins (Figure 16.2). In Africa, both diseases tend to be more severe at the top of the plant (Figures 16.1, 16.2).

Source and Transmission

Bacteria are spread within the field, from field to field and from infected weed hosts in wind-driven water droplets. Driving rains exacerbate the problem considerably. These diseases can also be seed transmitted. Debris from infected plants is a source of inoculum, as it infects overwintering weed hosts. In the semi-tropical areas where these diseases are a problem, winters are seldom cold enough to kill overwintering weeds.

Site Selection





THANK YOU

If you are interested – please contact me

anne.fisher@uky.edu